COLONOSCOPY PREPARATION

Five days prior to procedure:

Please STOP any NSAIDs which include Motrin, Ibuprofen, Aleve, and Advil.

Please **STOP** Flaxseed, Fish Oil and Biotin.

Three days prior to procedure:

START a low fiber diet:

AVOID seeds, nuts, corn, peas, beets or foods with skins on them such as tomatoes, cucumber, grapes, etc.

One day prior to procedure:

Begin clear liquid diet. You may have clear liquids ONLY.

Do not consume milk, alcohol, any red or purple colored liquids, or drinks that contain pulp.

STOP drinking liquids at least 4 hours before your colonoscopy.

Your procedure will be performed by Doctor: AHSCHUIER		
Address:		
Date of procedure:	Arrive at:	AM/PM
Comments:		

SUTAB

(sodium sulfate, magnesium sulfate, and potassium chloride) **Tablets**

1.479 g/0.225 g/0.188 g



On the Day Before Your Procedure

What You CAN Do

You may have clear liquids. ONLY

What You CANNOT Do

- Do not drink milk or eat or drink anything colored red or purple.
- Do not drink alcohol.
- Do not take other laxatives while taking SUTAB.
- Do not take oral medications within 1 hour of starting each dose of SUTAB.
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these medications at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB.

Liquids That Are **OK** to Drink

- Coffee or tea (no cream or nondairy creamer)
- Fruit juices (without pulp)
- Gelatin desserts (no fruit or topping)
- Water
- Chicken broth
- Clear soda (such as ginger ale)

NO red or purple

Note

- SUTAB is an osmotic laxative indicated for cleansing of the colon in preparation for colonoscopy in adults.
- Be sure to tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. SUTAB may affect how other medicines work.
- Medication taken by mouth may not be absorbed properly. when taken within 1 hour before the start of each dose of SUTAB.
- The most common adverse reactions after administration. of SUTAB were nausea, abdominal distension, vomiting, and upper abdominal pain.
- Contact your healthcare provider if you develop significant. vomiting or signs of dehydration after taking SUTAB or if you experience cardiac arrhythmias or seizures.
- If you have any questions about taking SUTAB, call your. doctor.

To learn more about this product, call 1-800-874-6756.

The Dosing Regimen

SUTAB is a split-dose (2-day) regimen. A total of 24 tablets is required for complete preparation for colonoscopy. You will take the tablets in two doses of 12 tablets each. Water must be consumed with each dose of SUTAB, and additional water must be consumed after each dose.


DOSE 1—On the Day Prior to Colonoscopy

Take the tablets with water

@ w:00 pm

STEP 1 Open 1 bottle of 12 tablets.

STEP 2 Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sipof water, and drink the entire amount of water over 15 to 20 minutes.



IMPORTANT: If you experience preparation-related symptoms (for example, nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.

Drink additional water

STEP 3 Approximately 1 hour after the last tablet is ingested, fill the provided container again with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.

STEP 4 Approximately 30 minutes after finishing the second container of water, fill the provided container with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.

DOSE 2—Day of the Colonoscopy

- Continue to consume only clear liquids until after the colonoscopy.

The morning of colonoscopy @ 2:00 AMI

open the second bottle of 12 tablets.

Repeat STEP 1 to STEP 4 from Dose 1.



IMPORTANT: You must complete all SUTAB tablets and required water at least a hours before colonoscopy.

Please read the full Prescribing Information and Medication Guide in the kit.

Provided as an educational service from 2 Braintree



©2020 Braintree Laboratories, Inc.

201-123-v1

December 2020